

Gliding dislocations in a driven vortex lattice

Stefan Scheidl¹ and Valerii M. Vinokur²

¹*Institut für Theoretische Physik, Universität zu Köln, Zùlpicher Str. 77, D-50937 Köln, Germany*

²*Materials Science Division, Argonne National Laboratory, Argonne, IL 60439*

(3 February 1997)

The dynamics of dislocations in a two-dimensional vortex lattice is studied in the presence of a pinning potential and a transport current. In a vortex lattice drifting with velocity v a glide velocity V_d of the dislocation with respect to the vortex lattice is found to decay like $V_d \sim v^{-4}$ for large drive. From this result the velocity for the crossover between a regime of coherent elastic motion and a regime of incoherent plastic motion of vortices is estimated.

PACS numbers: 74.60.Ge, 61.72.Bb

Periodic structures subject to quenched disorder are a generic model for a vast number of physical phenomena. The recent upraise of interest was motivated by the discovery of high temperature superconductors (HTS) that set a physical realization of the random system with tunable parameters and which are easily accessible in experiments [1]. As the statistical mechanics of the *static* periodic media in the presence of disorder reaches its mature phase, the physics of *non-equilibrium* processes proposes a diversity of unresolved challenging questions. In the static case it is established that quenched pinning destroys the crystalline order of the vortex lattice on intermediate spatial scales [1–3]. The long-range order of the vortex lattice is reduced to the quasi-long-range order of the “Bragg glass” [4,5]. It is important to note that in the static case and for weak pinning, the topological structure of the lattice is preserved in a certain range of the phase diagram [6].

One of the central issues of *non-equilibrium* statistical physics of dirty periodic media is the very nature of driven steady states. It was observed [7–9] that the depinning transition can be accompanied by plastic deformations of the driven periodic structures leading to the destruction of the moving solid. For driven vortex lattices it was proposed [9] that at larger drives a true dynamic phase transition occurs from an incoherently moving non-equilibrium state to a coherently moving solid phase. This conclusion was later extended to driven charge density waves [10]. Such dynamic transitions have indeed been observed in experiments [11] and simulations [9,12]. Note, that even in the limit of strong driving disorder severely affects the lattice structure on large length scales (in dimensions $d \leq 3$), which becomes strongly anisotropic and keeps glassy features [13–15].

A phenomenology of this dynamic melting is built on the notion that the driven periodic medium experiences pinning as temporally fluctuating distortion in the co-moving frame. A convenient heuristic tool to investigate the dynamic transition was introduced in Ref. [9], where the effect of temporarily fluctuating disorder was described as an additional thermal noise with an effective “shaking temperature”. This temperature increases

when the current is reduced and can lead to a “dynamic melting” of the lattice structure. While the phenomenological approach enables to estimate the position of the transition line on the phase diagram, the nature of the dynamic melting remains an open question. A microscopic understanding of dynamic melting requires the analysis of topological defects, which become free at the transition. In this Letter we present a first step in this direction focusing on the study the dynamics of *independent* dislocations in the driven two-dimensional vortex lattice. Note that in a *homogeneously* driven vortex lattice without pinning dislocations and vortices move with the same average velocity. Inhomogeneous pinning of different parts of the vortex lattice causes shear strains which generate a fluctuating force acting on dislocations. A motion of dislocation relative to the vortices arises from correlations of this force on *intermediate* scales. We examine the glide motion of the dislocations in the driven vortex lattice in the limit of large vortex drive, weak disorder, and well below the melting temperature of the pure vortex lattice. We calculate the dislocation velocity perturbatively and find that dislocations are retarded with respect to the vortices. The velocity where dislocations get pinned by disorder provides a non-equilibrium estimate for the crossover between elastic and plastic vortex flow.

In order to derive the equation of motion for the dislocation we first consider the dynamic response of the vortex lattice to disorder and thermal fluctuations. We restrict our discussion for simplicity to the two-dimensional vortex lattice. This lattice is supposed to move with an average velocity $\mathbf{v}$ as a consequence of the Lorentz force $\mathbf{f}_L$ generated by electric transport currents. We label vortices by their perfect lattice position $\mathbf{R}$ in the comoving frame. Their actual position in the laboratory frame is denoted by $\mathbf{r}$. Then a displacement can be defined as usual by $\mathbf{u}(\mathbf{R}, t) = \mathbf{r}(\mathbf{R}, t) - \mathbf{v}t$.

Neglecting nonlocal effects the elastic energy of the vortex lattice reads

$$\mathcal{H}_{\text{el}} = \frac{1}{2} \int \frac{d^2q}{(2\pi)^2} \{ c_{11} q^2 |\mathbf{u}^L(\mathbf{q})|^2 + c_{66} q^2 |\mathbf{u}^T(\mathbf{q})|^2 \}, \quad (1)$$

where $\mathbf{q}$ runs only over the first Brillouin zone and $\mathbf{u}^{L,T}$

are the longitudinal / transverse projections of the displacement field $\mathbf{u}(\mathbf{q}, \omega) = \int dt (\phi_0/B) \sum_{\mathbf{R}} e^{i\omega t - i\mathbf{q} \cdot \mathbf{R}} \mathbf{u}(\mathbf{R}, t)$ and ϕ_0/B is the area per vortex. The pinning potential $V(\mathbf{r})$ is supposed to have correlations $V(\mathbf{r})V(\mathbf{r}') = \Delta(\mathbf{r} - \mathbf{r}')$. The pinning energy of the lattice is $\mathcal{H}_{\text{pin}} = \sum_{\mathbf{R}} V(\mathbf{r}(\mathbf{R}))$.

The motion of vortices follows the overdamped Langevin equation

$$\eta \dot{\mathbf{r}}(\mathbf{R}, t) = -\frac{\partial}{\partial \mathbf{r}(\mathbf{R}, t)} [\mathcal{H}_{\text{el}} + \mathcal{H}_{\text{pin}}] + \mathbf{f}_L + \boldsymbol{\xi}(\mathbf{R}, t) \quad (2)$$

with the Bardeen-Stephen friction coefficient η and a thermal noise correlation $\langle \xi_\alpha(\mathbf{R}, t) \xi_\beta(\mathbf{R}', t') \rangle = 2\eta T \delta_{\alpha\beta} \delta_{\mathbf{R}\mathbf{R}'} \delta(t - t')$.

In the limit of large driving forces $\mathbf{f}_L$ the response of the vortices to the pinning potential vanishes. Then one has to leading order $\mathbf{v} = \mathbf{f}_L/\eta$ and the external force acting on the vortices is $\mathbf{f}(\mathbf{R}, t) := \boldsymbol{\xi}(\mathbf{R}, t) - \nabla V(\mathbf{R} + \mathbf{v}t)$. Its correlations read

$$\begin{aligned} \overline{\langle f_\alpha(\mathbf{q}, \omega) f_\beta(\mathbf{q}', \omega') \rangle} &= \Xi_{\alpha\beta}(\mathbf{q}, \omega) \delta(\mathbf{q} + \mathbf{q}') \delta(\omega + \omega'), \quad (3) \\ \Xi_{\alpha\beta}(\mathbf{q}, \omega) &= 2\eta T \frac{\phi_0}{B} \delta_{\alpha\beta} + \sum_{\mathbf{Q}} k_\alpha k_\beta \Delta(\mathbf{k}) \delta(\omega + \mathbf{k} \cdot \mathbf{v}), \end{aligned}$$

where $\mathbf{k} := \mathbf{Q} + \mathbf{q}$ with a reciprocal lattice vector $\mathbf{Q}$. In linear response [16] the vortex displacements are obtained from the forces $\mathbf{f}$ through the response function with longitudinal part $\Gamma^L(\mathbf{q}, \omega) = [-i\eta\omega + c_{11}(\phi_0/B)q^2]^{-1}$ and transverse part $\Gamma^T(\mathbf{q}, \omega) = [-i\eta\omega + c_{66}(\phi_0/B)q^2]^{-1}$. Typically the longitudinal response is small compared to the transverse response since $c_{11} \gg c_{66}$.

As one can immediately recognize from expression (3), pinning forces are more relevant than thermal forces [13] on *large* length and time scales: thermal fluctuations give a finite contribution to $\Xi_{\alpha\beta}(\mathbf{q} = 0, \omega = 0)$, whereas pinning forces give singular contributions for $\mathbf{Q} \neq 0$ with $\mathbf{Q} \cdot \mathbf{v} = 0$. This effect is due to the periodic structure of the vortex density and leads to the destruction of transitional long-range order for all velocities $\mathbf{v}$ in dimensions $d \leq 3$ [13,15]. For any finite driving the correct asymptotic behavior of the lattice roughness cannot be obtained in a linear response approach. However, as we show below, dislocation dynamics is governed by the elastic response of the vortex lattice on intermediate scales where this approach is sufficient.

The topological structure of the vortex lattice is expected to be destroyed by dislocations and disclinations, at least for sufficiently high temperature, strong disorder, and/or weak driving. Within an elastic approach vortices are found to move like in channels [13,14] when the lattice is moving along one of its principal directions. The displacement field component having the strongest fluctuations is the component parallel to the velocity [14,15]. This favours dislocations with Burgers vectors parallel to the velocity.

The motion of such a dislocation is easiest in the glide-direction parallel to the Burgers vector $\mathbf{b}$ with length of

the lattice spacing $a \approx \sqrt{\Phi_0/B}$. In this direction it can move by slips of individual vortices. A motion of the dislocation perpendicular to the Burgers vector (climb) is possible only in combination with the creation of vacancies or interstitials. In other terms, glide is controlled by the Peierls barrier which is much lower than the energy barriers for dislocation creep. In what follows we thus can ignore dislocation climb. The dislocation motion is then one-dimensional in the direction parallel to the Burgers vector and to the velocity. Its position along this direction in the *moving* vortex frame is denoted by $X(t)$, and we can consider $Y(t) = 0$ fixed.

According to the theory of dislocations [17] internal stresses

$$\sigma_{\alpha\beta} = 2c_{66} \nabla_\alpha u_\beta + (c_{11} - 2c_{66}) \nabla_\gamma u_\gamma \delta_{\alpha\beta} \quad (4)$$

of the lattice, due to thermal fluctuations and pinning, induce the so-called Peach-Koehler (PK) force

$$K_\alpha = \varepsilon_{\alpha\beta} \sigma_{\beta\gamma} b_\gamma \quad (5)$$

acting on the dislocation, where ε denotes the antisymmetric tensor. In addition, the dislocation is subject to a periodic potential $W(X) \approx -(W_p/2) \cos(2\pi X/a)$ with amplitude $W_p \approx 10^{-2} c_{66} a^2$ representing the Peierls barrier for glide motion. The resulting equation of motion for the dislocation reads

$$\eta_d \dot{X}(t) = K_x(X(t), t) - \nabla_x W(X(t)). \quad (6)$$

Here the dislocation friction coefficient is taken equal to its low velocity value $\eta_d \approx (\eta/8\pi) \ln(1/n_d a^2)$ [18,19] for an average dislocation density n_d . We assume that the vortex lattice is still ordered on intermediate length scales, i.e. the density of dislocations is small compared to the vortex density and $\eta_d \gg \eta$.

The fluctuating PK force includes thermal and disorder induced parts. The corresponding spatio-temporal correlator follows within linear response straightforwardly from Eq. (3) and has a rather complicated form. For the further analysis we single out the most important contributions.

One contribution arises from the thermal noise $\boldsymbol{\xi}$ acting on the vortices and it is present even in the absence of disorder and a driving force. It can be described as effective thermal noise $\xi_d(t)$ with correlations $\langle \xi_d(t) \xi_d(t') \rangle \approx 2\eta_d T \delta(t - t')$ experienced by *dislocations*.

The other contribution comes from pinning forces generating fluctuating displacements $\mathbf{u}(\mathbf{R}, t)$. As pointed out in Ref. [13], the modes $\mathbf{Q}$ with $\mathbf{Q} \cdot \mathbf{v} = 0$ of the force correlator (3) generate displacements, which are stationary in the laboratory frame, $\mathbf{u}^{(0)}(\mathbf{R}, t) = \mathbf{u}^{(0)}(\mathbf{R} + \mathbf{v}t)$. The associated contribution to the PK force is also stationary in the laboratory frame and can be the only origin of a pinning of dislocations by disorder. The correlations of these contributions $K_x^{(0)}$ at $Y = 0$ are

$$\overline{K_x^{(0)}(q_x)K_x^{(0)}(q'_x)} = \chi(q_x)\delta(q_x + q'_x), \quad (7)$$

$$\chi(q_x) \approx \frac{16a^2c_{66}^2\Delta_0}{3\pi a\eta^2v^2} \left(\frac{\eta v|q_x|}{c_{66}^2} \right)^{1/2}.$$

Here $\Delta_0 := \sum_{\mathbf{Q}: Q_x=0} Q_y^2 \Delta(\mathbf{Q})$ and the obtained formula is valid for $\eta v \gg c_{11}a$. Although the mode $\mathbf{Q} = 0$ also gives a stationary contribution, it is negligible as compared to the modes $\mathbf{Q} \neq 0$ with $\mathbf{Q} \cdot \mathbf{v} = 0$, because of $c_{11} \gg c_{66}$. We ignored the modes $\mathbf{Q} \cdot \mathbf{v} \neq 0$ which represent a shaking of the dislocation due to the discreteness of the vortex lattice and are expected only to renormalize the effective dislocation temperature. This effect is negligible for large $\mathbf{v}$, and does not contribute to the dislocation glide.

Having made the above approximations we rewrite the equation of motion (6) determining the dislocation glide velocity in a form

$$\eta_d \dot{X} = K_x^{(0)}(X + vt) - \frac{\pi W_p}{a} \sin \frac{2\pi X}{a} + \xi_d. \quad (8)$$

We restrict ourselves to temperatures $T \ll W_p$, i.e. well below the melting temperature $T_{KT} \approx (3c_{66}a^2)/8\pi$ of the pure lattice [20]. In this case the dislocation moves via thermal activation from one minimum of the Peierls potential to a neighboring one favored by the PK force. In the driven lattice disorder biases this motion in the direction opposite to the vortex drift.

For weak disorder $\Delta \ll T^2$ this effect is conveniently calculated perturbatively in the dynamical functional formalism [21], where the dislocation position X and its response field $\tilde{X}$ are assigned a statistical weight $\exp(-\mathcal{A})$ with a dynamic action

$$\mathcal{A} = \mathcal{A}_0 + \mathcal{A}_1, \quad (9)$$

$$\mathcal{A}_0 = \int dt \{ \eta_d T \tilde{X}^2 + i\tilde{X}[\eta_d \dot{X} + \nabla_X W(X)] \},$$

$$\mathcal{A}_1 = \frac{1}{2} \int dt \int dt' \tilde{X} \tilde{X}' \chi(X - X' + v(t - t')).$$

Here the disorder average is already performed and abbreviations $X \equiv X(t)$, $X' \equiv X(t')$ etc. are used.

In the absence of disorder, dynamics is controlled by the free action $\mathcal{A}_0$. This action is minimized by instanton trajectories which start from a minimum/maximum of W at $t = -\infty$ and end at a neighbored maximum/minimum at $t = \infty$:

$$X_i(t) = \sigma_2 \frac{a}{\pi} \arctan \left[\exp \left(\sigma_1 \frac{t}{t_i} \right) \right], \quad (10a)$$

$$\tilde{X}_i(t) = -\frac{i}{2\eta_d T} [\eta_d \dot{X}_i(t) + \nabla_X W(X_i)], \quad (10b)$$

where $t_i = a^2\eta_d/2\pi^2W_p$ is an instanton time scale and $\sigma_1, \sigma_2 = \pm 1$ specifies the type. For $\sigma_1 = \pm 1$ motion is up or down from a minimum to a maximum of W or vice versa. For $\sigma_1\sigma_2 = \pm 1$ motion is forward/backward in X

direction. Both “down”-instantons have $\tilde{X}_i^{\text{down}}(t) = 0$ and thus $\mathcal{A}_0^{\text{down}} = 0$. Both “up”-instantons have

$$\mathcal{A}_0^{\text{up}} = W_p/T, \quad (11a)$$

$$\tilde{X}_i^{\text{up}}(t) = -\frac{i\sigma_2 a}{2\pi t_i T} \frac{1}{\cosh(t/t_i)}. \quad (11b)$$

Eq. (11a) gives the usual statistical weight of thermally activated instantons.

Now one can easily calculate the disorder corrections to the instanton actions in perturbation theory. To evaluate $\mathcal{A}_1$ in the lowest order in Δ one has to use the *unperturbed* instanton solutions and recognizes immediately that $\mathcal{A}_1^{\text{down}} = 0$ since $\tilde{X}^{\text{down}} = 0$. Near its saddle-point the dislocation has a velocity $v_i = v + \dot{X}_i(0)$ relative to the disorder, which is different for “forward” and “backward” instantons. In terms of this velocity the action correction reads

$$\mathcal{A}_1 \approx \frac{1}{2} \int dt \int dt' \tilde{X} \tilde{X}' \chi(v_i(t - t'))$$

$$\approx -\frac{2\Delta_0}{3\pi^2 T^2} \left(\frac{2a^2 c_{66}}{\pi \eta v v_i t_i} \right)^{3/2}. \quad (12)$$

The resulting dislocation glide velocity $V_d := \langle \dot{X} \rangle$ is obtained from the difference of activation rates for forward and backward instantons, $V_d \approx (a/t_0)[\exp(-\mathcal{A}^{\text{fw,up}}) - \exp(-\mathcal{A}^{\text{bw,up}})]$. The attempt frequency of the dislocation $t_0 \approx a^2\eta_d/\pi W_p$ is estimated according to Kramers rate theory [22]. In terms of the dislocation diffusion constant $D_0 = (a^2/t_0) \exp(-\mathcal{A}_0)$ in the pure lattice we eventually find a drift velocity

$$V_d \approx -\frac{D_0\Delta_0}{vt_i T^2} \left(\frac{2a^2 c_{66}}{\pi^3 \eta v^2 t_i} \right)^{3/2} \propto v^{-4} \quad (13)$$

in the limit $v \rightarrow \infty$, where $\mathcal{A}_1 \ll 1$.

Several properties of this resulting drift velocity are noteworthy: it decays with the fourth power of velocity, which is much faster than $V_d \sim v^{-1}$ that one might expect on the basis of symmetry arguments. The drift velocity is independent of c_{11} for $c_{11} \gg c_{66}$, where only shear modes of the lattice get excited. In the considered temperature range $T \ll W_p$ the drift velocity arises from a small bias of instantonic hops. Since this process is thermally activated the drift velocity increases monotonously with temperature. The asymptotic form (13) holds only down to a certain minimum temperature determined by the condition $\mathcal{A}_1 \ll 1$, which however is satisfied for large velocity.

From Eq. (13) we can estimate the critical velocity where a crossover from “elastic” to “plastic” flow takes place. In the “elastic” flow regime vortices flow essentially coherently. Topological defects may be present due to thermal fluctuations. Although we found that dislocations are always retarded with respect to the vortices, this effect is small for high drift velocities $v \gg |V_d|$. On the other hand there is the “plastic” flow regime, where

dislocations are essentially pinned by impurities. Such dislocations are a source of rearrangements of neighborhoods between vortices. This plasticity becomes significant for $1 - |V_d|/v \ll 1$. Although our derivation of Eq. (13) is strictly valid only for large velocities, we expect it to give a qualitatively correct estimate for the velocity

$$v_c \approx \left[\frac{D_0 \Delta_0}{t_i T^2} \left(\frac{2a^2 c_{66}}{\pi^3 \eta t_i} \right)^{3/2} \right]^{1/5} \quad (14)$$

of the crossover between both regimes.

Although the instantaneous forces acting on the dislocation arise from local stresses in the vortex lattice, the dislocation glide velocity results from the correlations of these stresses on *intermediate* scales $|X| \lesssim vt_i$ for large drift velocities: According to Eq. (11b) the quantity $\tilde{X}_i(t)$ decays exponentially on the instanton time scale t_i . Thus the main contributions to $\mathcal{A}_1$ in Eq. (9) come from $\chi(X)$ for $|X| \lesssim vt_i$. This scale increases $\propto v^1$, whereas the dynamic Larkin length, up to where the perturbative calculation of χ is reliable, increases $\propto v^3$ [13,15]. It is for this reason that the calculation of χ in linear response approximation for the vortex lattice is sufficient for the dislocation dynamics at large drive.

Relating our result $v_c \propto \Delta_0^{1/5}$ with the finding [9] of a dynamic freezing velocity $v_f \propto \Delta_0$, one has to distinguish the case of weak and strong pinning. For *strong* pinning $v_c < v_f$. Expression (14) for v_c is then formally invalid since the lattice is no longer ordered at these velocities. In this case we can describe only the regime $v > v_f$, where the vortex lattice preserves its crystalline order on large scales and dislocations move with the vortices. In the limit of *weak* pinning both velocities $v_f < v_c$ become small and the perturbative findings are no longer quantitatively reliable. Nevertheless, the smallness of the velocities indicates that the lattice is well ordered and dislocations move essentially with the lattice at practically all drift velocities. Qualitatively, our finding suggests a crossover between practically elastic flow for $v > v_c$ and strongly plastic flow for $v < v_c$. This crossover can be viewed as precursor of the pinning transition.

To sum up, our result $V_d \propto v^{-4}$ implies that the motion of the vortex lattice in disorder is *always* accompanied by plasticity, even if it is extremely weak at high drives. For weak pinning we have estimated the characteristic velocity of the crossover between “elastic” and “plastic” flow of the vortex lattice. We expect these results to be valid also after the inclusion of interactions between dislocations. For example a dislocation-antidislocation pair will be retarded with respect to the vortex lattice essentially as a whole, since the resulting glide velocity is independent of the dislocation type. But even if this pair remains bound it can create point defects during its motion. Interactions between dislocations are clearly relevant for a study of the qualitative nature of the dynamic melting

transition [9], we will address this issue in a forthcoming publication.

The authors gratefully acknowledge discussions with G.T. Zimányi. This work was supported from Argonne National Laboratory through the U.S. Department of Energy, BES-Material Sciences, under contract No. W-31-109-ENG-38 and by the NSF-Office of Science and Technology Centers under contract No. DMR91-20000 Science and Technology Center for Superconductivity and by Deutsche Forschungsgemeinschaft project SFB341.

-
- [1] G. Blatter *et al.*, Rev. Mod. Phys. **66**, 1125 (1994).
 - [2] A. I. Larkin, Sov. Phys. JETP **31**, 784 (1970).
 - [3] M. V. Feigelman *et al.*, Phys. Rev. Lett. **63**, 2303, (1989).
 - [4] T. Nattermann, Phys. Rev. Lett. **64**, 2454 (1990).
 - [5] T. Giamarchi and P. Le Doussal, Phys. Rev. Lett. **72**, 1530 (1994); Phys. Rev. B **52**, 1242 (1995).
 - [6] J. Kierfeld, T. Nattermann, and T. Hwa, preprint cond-mat/9512101; J. Kierfeld, preprint cond-mat/9609045; D. Ertas and D.R. Nelson, preprint cond-mat/9607142; D. Carpentier, P. Le Doussal, and T. Giamarchi, Europhys. Lett. **35**, 379 (1996); T. Giamarchi and P. Le Doussal, preprint cond-mat/9609112.
 - [7] S. N. Coppersmith, Phys. Rev. Lett., B **44**, 2887 (1991).
 - [8] H. J. Jensen *et al.*, J. Low Temp. Phys. **74**, 293 (1989).
 - [9] A.E. Koshelev and V.M. Vinokur, Phys. Rev. Lett. **73**, 3580 (1994).
 - [10] L. Balents and M.P.A. Fisher, Phys. Rev. Lett. **75**, 4270 (1995).
 - [11] M.C. Hellerqvist *et al.*, Phys. Rev. Lett. **76**, 4022 (1996).
 - [12] S. Ryu *et al.*, preprint cond-mat/9606126.
 - [13] T. Giamarchi and P. Le Doussal, Phys. Rev. Lett. **76**, 3408 (1996); preprint cond-mat/9608153.
 - [14] K. Moon, R.T. Scalettar, and G.T. Zimányi, Phys. Rev. Lett. **77**, 2778 (1996).
 - [15] L. Balents, M.C. Marchetti, and L. Radzihovski, preprint cond-mat/9608009.
 - [16] A. Schmid and W. Hauger, J. Low Temp. Phys. **11**, 667 (1973).
 - [17] see, e.g., J.P. Hirth and J. Lothe, “Theory of dislocations” (Wiley & Sons, New York, 1982).
 - [18] E.H. Brandt, Phys. Stat. Sol. **36**, K167 (1969).
 - [19] A.E. Koshelev, Phys. Rev. B **45**, 12936 (1992).
 - [20] J.M. Kosterlitz and D.J. Thouless, J. Phys. C **6**, 1181 (1973).
 - [21] P.C. Martin, E.D. Siggia, and H.A. Rose, Phys. Rev. A **8**, 423 (1973).
 - [22] P. Hänggi, P. Talkner, and M. Borkovec, Rev. Mod. Phys. **62**, 251 (1990).